Physical Activity

Health benefits to your get-up-and-go

Types of Physical Activity

Aerobic - moving large muscles in a rhythmic way to get your heart pumping

Strength - muscle contractions against resistance

Flexibility - the range of motion possible at a joint

Balance - maintaining body's equilibrium while stationary or moving

Benefits

Aerobic	lowers risk of all-cause mortality	increased quality of life
	lower incidence of cancer (multiple types)	increased quality of sleep
	lowers incidence & mortality cardiovascular d	isease much, much more!
Strength	Reduced injury	Improved quality of life
	Increased basal metabolic rate	Improved quality of sleep
	Reduced fatigue Improved bone mineral density post-menopause	
	Improved sleep	Improved weight maintenance
	Improved performance and muscle strength	Decreased pain/disability
	Reduced risk of falls in the elderly	Improved mental health
Flexibility	Improved recovery/healing	Reduces risk of injury
	Improved recovery (soreness)	Enhances physical performance
Balance	Improves body's ability to withstand destabilizing stimuli or postural sway	
	Decreases injury in elderly population	
	37% decrease in injuries from falls	
	30% decrease in injuries requiring medical care	
	43% decrease in severe injury falls	
	61% decrease in falls resulting in fractures	

Health Research

If worldwide physical activity increased by 25%, then 1.3M deaths could be averted each year (1).

Sedentary activity (no movement) is especially to blame; causes 7% mortality (all causes) (2)

Standing for 2 hrs per day (as opposed to sitting) will reduce all-cause mortality by 10% (3), BUT physical activity of just 1-1.9 hrs moderate to vigorous activity (able to talk but not sing) per week will reduce it by 18% (4)

Moderate physical activity reduces risk of coronary heart disease in active persons by 20-30% when compared to inactive persons (5)

Physical activity decreases risk of Type 2 Diabetes by 31% (6) AND was effective to lower HbA1C levels with or without diet co-intervention (7)

Physical activity decreases risk of cancers (especially breast & colon), likely at about 3 hrs per week of moderate to vigorous intensity exercise (8). Colon cancer risk is decreased by 17-30% with exercise (9)

Years of Life gained (10):

- 1. 75 mins brisk walking per week = ~1.8 years
- 2. 90-150 mins brisk walking per week = ~2.5 years
- 3. 200 mins brisk walking per week = ~3 yrs
- 4. 400 mins brisk walking per week = ~4 yrs
- 5. 500 mins brisk walking per week = ~4.5 yrs

References

- 1. Lee IM et al. Effect of physical activity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012:380(9838):219-229. http://doi.org/10.1016/s0140-6736(12)61031-9
- 2. van der Ploeg HP, et al. Sitting time and all-cause mortality risk in 222,497 Australian adults. Arch Intern Med. 2012:172(6):494-500. http://doi.org/10.1001/archinternmed.2011.2174
- 3. van der Ploeg HP, et al. Standing time and all-cause mortality in a large cohort of Australian adults. *Prev Med.* 2014;69:187-191. http://doi.org/10.1016/j.ypmed.201410.004
- 4. Wen CP, et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet*. 2011:378(9798):1244-1253. http://doi.org/10.1016/s0140-6736(11)60749-6
- 5. Darden D, et al. Physical Activity and Exercise for Secondary Prevention among Patients with Cardiovascular Disease. *Curr Cardiovasc Risk Rep.* 2013;7(6), http://doi.org/10.1007/s12170-013-0354-5
- 6. Jeon CY, et al. Physical activity of moderate intensity and risk of type 2 diabetes: a systematic review. *Diabetes Care*. 2007;30(3):744-752. http://doi.org//10.2337/dc06-1842
- 7. Umpierre D, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 2011;305(17):1790-1799. http://doi.org/10.1001/jama.2011.576
- 8. Kushi LH, et al. American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin.* 2012;62(1):30-67
- 9. Wolin KY, et al. Physical activity and colon cancer prevention: a meta-analysis. *Br J Cancer.* 2009;100(4);611-616. http://doi.org/10.1038/sj.bjc.6604917
- 10. Woodcock J, et al. Non-vigorous physical activity and all-cause mortality: systematic review and meta-analysis of cohort studies. *Int J Epidemiol*. 2011:40(1):121-138. http://doi.org/10.1093/ije/dyq104

Information adapted from Clayton JS. Foundations of Lifestyle Medicine Board Review Manual, 4th ed. Section 6: Physical Activity Science and Prescription. American College of Lifestyle Medicine.